

# SANITEK SANITIZE

**DISINFECTANT • SANITIZER  
DETERGENT • DEODORIZER**

**DESCRIPTION:**

SANITEK SANITIZE is a phosphate-free formulation designed to provide effective cleaning, deodorizing, and disinfection specifically for hospitals, nursing homes, schools, food processing plants, food service establishments, and other institutions where housekeeping is of prime importance in controlling the hazard of cross contamination.

**EPA REGISTERED:**

EPA REG. NO. 1839-79-2839 EPA EST NO. 2839-CA-1

**INGREDIENTS:**

**ACTIVE INGREDIENTS:** n-Alkyl (60% C<sub>14</sub>, 30% C<sub>16</sub>, 5% C<sub>12</sub>, 5% C<sub>18</sub>)  
dimethyl benzyl ammonium chlorides.....2.25%  
n-Alkyl (68% C<sub>12</sub>, 32% C<sub>14</sub>)  
dimethyl ethylbenzyl ammonium chlorides .....2.25%  
**INERT INGREDIENTS:** .....95.50%  
**TOTAL** .....100.00%

**U.S.D.A. REGISTERED:**

Category D-2. When used for sanitation of previously cleaned food equipment or food contact items, limit active quaternaries to 200 ppm. At this level NO POTABLE WATER RINSE IS REQUIRED. SANITEK SANITIZE is an effective sanitizer when diluted in water up to 250 ppm hardness (as CaCO<sub>3</sub>).

**DIRECTIONS FOR USE:**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

**SANITIZATION** - To sanitize previously cleaned and rinsed non-porous food contact surfaces, prepare a 200 ppm active quaternary solution by adding 2 ounces SANITEK SANITIZE to 3.5 gallons of water.

To sanitize immobile items such as tanks, chopping blocks, and counter tops, flood the area with a 200 ppm active quaternary solution or apply with a cloth or sponge making sure to wet all surfaces completely for at least 60 seconds. Let air dry. Prepare a fresh solution for each use.

To sanitize mobile items such as drinking glasses and eating utensils, immerse in a 200 ppm active quaternary solution for at least 60 seconds making sure to immerse completely. Remove and let air dry. Prepare a fresh solution daily or more frequently as soil is apparent.

**DISINFECTION** - To disinfect inanimate hard surfaces, add 2 ounces SANITEK SANITIZE per gallon of water. Apply solution with a mop, cloth, sponge, or mechanical sprayer so as to wet all surfaces thoroughly. Allow to remain wet for 10 minutes, then remove excess liquid. For heavily soiled areas, a pre-cleaning step is required; prepare a fresh solution for each use. To disinfect toilet bowls, flush toilet and add 2 ounces SANITEK SANITIZE directly to the bowl water. Swab the bowl completely using a scrub brush or toilet mop, making sure to get under the rim. Let stand for 10 minutes, then flush.

**EFFICACY TESTS HAVE DEMONSTRATED THAT THIS PRODUCT IS AN EFFECTIVE BACTERICIDE, FUNGICIDE, AND VIRUCIDE, IN THE PRESENCE OF ORGANIC SOIL (5% BLOOD SERUM).**

**BACTERICIDAL ACTIVITY** - When diluted at the rate of 2 ounces per gallon of water, SANITEK SANITIZE exhibits effective disinfectant activity against the organisms: Pseudomonas aeruginosa PRD 10, Escherichia coli, Klebsiella pneumoniae, Salmonella schottmuelleri, Salmonella choleraesuis, Streptococcus salivarius, Brevibacterium ammoniagenes, Streptococcus faecalis, Shigella dysenteriae, Enterobacter aerogenes, and Staphylococcus aureus, and meets all requirements for hospital use.

**SANITEK SANITIZE KILLS HIV ON PRE-CLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS** in health care settings (Hospitals, Nursing Homes) or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of human immunodeficiency virus Type 1 (HIV-1) (associated with AIDS.)

**SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 OF SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS. PERSONAL PROTECTION:** When handling items soiled with blood or body fluids, use disposable latex gloves, gowns, masks, or eye coverings. **CLEANING PROCEDURES:** Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of SANITEK SANITIZE. **CONTACT TIME:** Allow surface to remain wet for 10 minutes. **DISPOSAL OF INFECTIOUS MATERIALS:** Blood and other body fluids should be autoclaved and disposed of according to local regulations for infectious waste disposal.

**HEALTH & SAFETY:**

**KEEP OUT OF REACH OF CHILDREN  
DANGER**

See Label and MSDS for More Health & Safety Information.

**PACKAGING:**

12/1-pint case, 4/1-gallon case, 5-gallon pails, 55-gallon drums

**ITEM NO.:**

330

04/98



**SANITEK**

CHEMICAL SPECIALTIES MANUFACTURER

Sanitek Products, Inc. • 3959 Goodwin Ave., Los Angeles, CA 90039 • Phone (323) 245-6781